

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, C12N 15/00	A1	(11) International Publication Number: WO 99/18239 (43) International Publication Date: 15 April 1999 (15.04.99)
(21) International Application Number: PCT/US98/20582 (22) International Filing Date: 1 October 1998 (01.10.98) (30) Priority Data: 60/060,767 2 October 1997 (02.10.97) US (71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BLACK, Michael, Terence [GB/FR]; 47, Boulevard d'Angleterre, F-78110 Le Vesinet (FR). KALLENDER, Howard [GB/US]; 1220 Bill Smith Boulevard, King of Prussia, PA 19406 (US). (74) Agents: GIMMI, Edward, R. et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: ANTIMICROBIAL DRUG SCREENING USING A RECOMBINANT CELL COMPRISING A RNA-DEPENDENT AMIDOTRANSFERASE GENE (57) Abstract This invention relates to newly developed methods for discovering antimicrobial compounds using an RAT-based assay system. This invention also relates to compositions of matter useful in carrying out the methods of the invention as well as antimicrobial compounds developed using such methods.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

ANTIMICROBIAL DRUG SCREENING USING A RECOMBINANT CELL COMPRISING A RNA-DEPENDENT AMIDOTRANSFERASE GENE

FIELD OF THE INVENTION

This invention relates to newly developed methods for discovering antimicrobial compounds using a RAT gene-based whole cell assay. It is particularly suited for carrying out antimicrobial compound screening assays in bacterial cells. This invention also relates to compositions of matter useful in carrying out the methods of the invention as well as antimicrobial compounds developed using such methods.

BACKGROUND OF THE INVENTION

Despite the absence of glutamyl-tRNA synthetase activity in Gram-positive bacteria, and likely all Gram-negative bacteria with the exception of the gamma subdivision of purple bacteria, bacterial cells are clearly able to produce the Gln-tRNA(Gln) required for accurate protein synthesis. The mechanism by which this is achieved involves the formation of Glu-tRNA(Gln) as an intermediate that is produced by the misaminoacylation of tRNA(Gln) by glutamyl-tRNA synthetase (ERS). This reaction would be toxic as it would lead to Gln-tRNA(Gln) starvation and to the synthesis of aberrant proteins and the consequent cessation of bacterial protein synthesis. However, the 'correct' end product, Gln-tRNA(Gln), is formed from Glu-tRNA(Gln) by transfer of an amine group to the ligated glutamate residue. This reaction is catalyzed by a tRNA- and Mg^{2+} /ATP-dependent amidotransferase. (RNA-dependent AmidoTransferase - RAT) ; also known as Glu-tRNA^{Gln} amidotransferase or Glu-AdT - Curnow-AW, *et al. PNAS* 94, 11819-11826 (1997). Inhibition of this apparently ubiquitous reaction in Gram-positive organisms, and some Gram-negative organisms, would effectively lead to Gln-tRNA(Gln) starvation and to cell death.

This invention provides methods for exploiting the relationship between RAT and glutamyl tRNA synthetase to screen for antimicrobial compounds that target either or both of these enzymes, for example by binding to them or affecting their enzymatic pathways. There is a need for methods for screening for novel antimicrobial compounds, such as the screening methods of the invention. Such methods have a present benefit of being useful to screen compounds for antibiotic activity that can play a role in preventing, ameliorating or correcting infections, dysfunctions or diseases, such as bacterial infections.

GLOSSARY

The following definitions are provided to facilitate understanding of certain terms used frequently herein. Certain other definitions are provided elsewhere herein.

"Host cell" is a cell which has been transformed or transfected, or is capable of transformation or transfection by an exogenous polynucleotide sequence.

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein.

"Polynucleotide(s)" generally refers to any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotide(s)" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions or single-, double- and triple-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded, or triple-stranded regions, or a mixture of single- and double-stranded regions. In addition, "polynucleotide" as used herein refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The strands in such regions may be from the same molecule or from different molecules. The regions may include all of one or more of the molecules, but more typically involve only a region of some of the molecules. One of the molecules of a triple-helical region often is an oligonucleotide. As used herein, the term "polynucleotide(s)" also includes DNAs or RNAs as described above that contain one or more modified bases. Thus, DNAs or RNAs with backbones modified for stability or for other reasons are "polynucleotide(s)" as that term is intended herein. Moreover, DNAs or RNAs comprising unusual bases, such as inosine, or modified bases, such as tritylated bases, to name just two examples, are polynucleotides as the term is used herein. It will be appreciated that a great variety of modifications have been made to DNA and RNA that serve many useful purposes known to those of skill in the art. The term "polynucleotide(s)" as it is employed herein embraces such chemically, enzymatically or metabolically modified forms of polynucleotides, as well as the chemical forms of DNA and RNA characteristic of viruses and cells, including, for example, simple and complex cells. "Polynucleotide(s)" also embraces short polynucleotides often referred to as oligonucleotide(s).

"Polypeptide(s)" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds. "Polypeptide(s)" refers to both short chains, commonly referred to as peptides, oligopeptides and oligomers and to longer chains generally referred to as proteins. Polypeptides may contain amino acids other

than the 20 gene encoded amino acids. "Polypeptide(s)" include those modified either by natural processes, such as processing and other post-translational modifications, but also by chemical modification techniques. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature, and they are well known to those of skill in the art. It will be appreciated that the same type of modification may be present in the same or varying degree at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains, and the amino or carboxyl termini. Modifications include, for example, acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, glycosylation, lipid attachment, sulfation, gamma-carboxylation of glutamic acid residues, hydroxylation and ADP-ribosylation, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins, such as arginylation, and ubiquitination. See, for instance, *PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993) and Wold, F., *Posttranslational Protein Modifications: Perspectives and Prospects*, pgs. 1-12 in *POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS*, B. C. Johnson, Ed., Academic Press, New York (1983); Seifter et al., *Meth. Enzymol.* 182:626-646 (1990) and Rattan et al., *Protein Synthesis: Posttranslational Modifications and Aging*, Ann. N.Y. Acad. Sci. 663: 48-62 (1992). Polypeptides may be branched or cyclic, with or without branching. Cyclic, branched and branched circular polypeptides may result from post-translational natural processes and may be made by entirely synthetic methods, as well.

"Variant(s)" as the term is used herein, is a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide respectively, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions,

deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques, by direct synthesis, and by other recombinant methods known to skilled artisans.

SUMMARY OF THE INVENTION

A method of screening for antimicrobial drugs comprising the steps of: providing at least one cell naturally lacking a RAT gene or genes and comprising at least one recombinant glutamyl tRNA synthetase gene and at least one recombinant RAT gene; contacting the cell with at least one candidate compound; and detecting altered metabolism in the cell of the contacting step. Methods are also provided for screening in *Staphylococcus aureus* that uses a wild type glutamyl tRNA synthetase gene.

A method wherein the recombinant glutamyl tRNA synthetase and RAT genes are on an episomal element or integrated into a chromosome of the cell.

A method wherein RAT gene expression level is regulated.

A method wherein a RAT gene is a RAT gene selected from the group consisting of a Gram positive bacteria, a Gram negative bacteria, a streptococcus, *S. pneumoniae*, a staphylococcus, *S. aureus*, enterococci, *Enterococcus faecalis*, *Enterococcus faecium*, a *Bacillus*, and *Bacillus subtilis*.

A method wherein the glutamyl tRNA synthetase gene is a glutamyl tRNA synthetase gene selected from the group consisting of a Gram positive bacteria, a Gram negative bacteria, a streptococcus, *S. pneumoniae*, a staphylococcus, *S. aureus*, enterococci, *Enterococcus faecalis*, *Enterococcus faecium*, a *Bacillus*, and *Bacillus subtilis*.

A method wherein the altered metabolism comprises inhibition of RAT protein activity.

A method wherein the detecting step further comprises detecting, an accumulation, particularly a toxic accumulation, of Glu-tRNA(Gln) or a toxic incorporation of a glutamyl residue for a glutaminy in a nascent protein chain.

5 A method wherein the detecting step further comprises detecting cell death or a reduction in growth rate or amount, particularly a substantial reduction in growth rate or amount.

A method wherein the cell possesses a glutaminy tRNA synthetase or lacks a RAT gene.

10 A method wherein the cell lacks a glutaminy tRNA synthetase or possesses a RAT gene, particularly *S. aureus*.

An isolated bacterial cell lacking a RAT gene and comprising at least one recombinant bacterial glutamyl tRNA synthetase gene and at least one recombinant bacterial RAT gene.

15 An isolated bacterial cell possessing a RAT gene and comprising at least one recombinant or wild type bacterial glutamyl tRNA synthetase gene and at least one recombinant bacterial RAT gene.

A method or composition wherein the glutamyl tRNA synthetase and RAT genes are on episomal element or integrated into a chromosome of the cell.

A host cell wherein the RAT gene expression level is regulated.

20 A host cell wherein a RAT gene is an *S. aureus* RAT gene.

A host cell wherein the glutamyl tRNA synthetase gene is a *S. aureus* glutamyl tRNA synthetase gene.

A host cell wherein the cell possesses a glutaminy tRNA synthetase or lacks a RAT gene.

25 A host cell wherein the cell lacks a glutaminy tRNA synthetase or possesses a RAT gene.

A method wherein the detecting step further comprises detecting altered translation.

30 A method wherein the detecting step further comprises detecting altered test protein.

A kit comprising at least one bacterial cell lacking a RAT gene and the cell comprising at least one bacterial glutamyl tRNA synthetase gene and at least one bacterial RAT gene.

A kit comprising a polynucleotide encoding a RAT gene and a polynucleotide encoding a glutamyl tRNA synthetase gene.

A polynucleotide comprising a RAT expressibly linked to an inducible promoter.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Figures 1 shows a schematic diagram of a preferred embodiment of the invention showing a schematic diagram of the putative mechanism of an embodiment of the invention.

Figure 2 shows a graph demonstrating the putative mechanism of one preferred embodiment of the invention.

10 Figures 3 shows a schematic diagram of a preferred embodiment of the invention showing a schematic diagram of the putative mechanism of an embodiment of the invention.

DETAILED DESCRIPTION OF THE INVENTION

15 The present invention provides a method for screening for compounds that lead to or are involved in aborted translation, particularly those that do not inhibit protein synthesis *per se*. Applicants believed that such compounds will have an enormous impact on 'global' protein synthesis, and are therefore predicted to be bactericidal.

20 The natural promoter of the RAT gene (herein "RAT gene(s)" means a gene encoding any or all of the RAT protein subunits in an enzymatically active or biologically functional RAT protein complex (e.g., a RAT gene in Table 1), and "RAT protein(s)" means any protein encoded by a RAT gene (e.g., a RAT protein in Table 1)) will be replaced with a heterologous, regulatable promoter (e.g., an inducible promoter) in the chromosome of a RAT-expressing bacterial host cell, such as by homologous recombination (in a preferred embodiment insertional mutagenesis is used since, for
25 example, it is more rapid than a double crossover and should give the same phenotype. Such RAT gene constructs comprising a regulatable promoter is referred to as "hybrid RAT." However, by using this method there will be an extra copy of the first stretch of base pairs (e.g., about 300-700 base pairs, preferably about 500 base pairs) of the RAT gene present, still under the control of the native RAT promoter. This will not be sufficient
30 sequence to encode active RAT).

Preferred host cells useful in the invention include, but are not limited to, any bacteria comprising a natural endogenous RAT gene, such as, any Gram positive bacteria, many Gram negative bacteria, and also a member of the genus *Streptococcus*,

- Staphylococcus*, *Bordetella*, *Corynebacterium*, *Mycobacterium*, *Neisseria*, *Haemophilus*, *Actinomycetes*, *Streptomyces*, *Nocardia*, *Enterobacter*, *Yersinia*, *Fancisella*, *Pasturella*, *Moraxella*, *Acinetobacter*, *Erysipelothrix*, *Branhamella*, *Actinobacillus*, *Streptobacillus*, *Listeria*, *Calymmatobacterium*, *Brucella*, *Bacillus*, *Clostridium*, *Treponema*, *Escherichia*,
5 *Salmonella*, *Kleibsiella*, *Vibrio*, *Proteus*, *Erwinia*, *Borrelia*, *Leptospira*, *Spirillum*, *Campylobacter*, *Shigella*, *Legionella*, *Pseudomonas*, *Aeromonas*, *Rickettsia*, *Chlamydia*, *Borrelia* and *Mycoplasma*, and further including, but not limited to, a member of the species or group, Group A *Streptococcus*, Group B *Streptococcus*, Group C *Streptococcus*, Group D *Streptococcus*, Group G *Streptococcus*, *Streptococcus pneumoniae*, *Streptococcus pyogenes*,
10 *Streptococcus agalactiae*, *Streptococcus faecalis*, *Streptococcus faecium*, *Streptococcus durans*, *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Staphylococcus aureus*, particularly *Staphylococcus aureus* strain RN4220, *Staphylococcus epidermidis*, *Corynebacterium diphtheriae*, *Gardnerella vaginalis*, *Mycobacterium tuberculosis*, *Mycobacterium bovis*, *Mycobacterium ulcerans*, *Mycobacterium leprae*, *Actinomycetes israelii*, *Listeria monocytogenes*, *Bordetella pertussis*, *Bordetella parapertussis*, *Bordetella bronchiseptica*,
15 *Escherichia coli*, *Shigella dysenteriae*, *Haemophilus influenzae*, *Haemophilus aegyptius*, *Haemophilus parainfluenzae*, *Haemophilus ducreyi*, *Bordetella*, *Salmonella typhi*, *Citrobacter freundii*, *Proteus mirabilis*, *Proteus vulgaris*, *Yersinia pestis*, *Kleibsiella pneumoniae*, *Serratia marcescens*, *Serratia liquefaciens*, *Vibrio cholera*, *Shigella dysenterii*, *Shigella flexneri*, *Pseudomonas aeruginosa*, *Francisella tularensis*, *Brucella abortus*, *Bacillus anthracis*, *Bacillus cereus*, *Clostridium perfringens*, *Clostridium tetani*, *Clostridium botulinum*, *Treponema pallidum*, *Rickettsia rickettsii* and *Chlamydia trachomatis*, (ii) an archaeon, including but not limited to *Archaeobacter*, and (iii) a unicellular or filamentous eukaryote, including but not limited to, a protozoan, a fungus, a member of the genus
25 *Saccharomyces*, *Kluveromyces*, or *Candida*, and a member of the species *Saccharomyces cerevisiae*, *Kluveromyces lactis*, or *Candida albicans*.

Regulatable promoters, particularly inducible promoters useful in the invention include, but are not limited to, P_{xylA} plus the *xylR* repressor gene, from various bacteria, such as *Bacillus* sp. and *Lactobacillus pentosus*; P_{lacA} plus the *lacR* repressor gene, from
30 various bacteria, such as *E. coli* and *Lactococcus lacti*; hybrid promoters consisting of the *E. coli lac* repressor/operator and the -10 and -35 regions of various promoters, such as phages SP0-1 (known as P_{spac}) and T5; $P_{xyl/tet}$ - a hybrid consisting of the *E. coli Tn10 tet* repressor/operator and the *Bacillus subtilis xylA* -10 and -35 regions; P_{T7} plus the T7 RNA

polymerase gene under the control of one of the described promoters; P_{trp} from various bacteria; $\phi 31$ middle promoter from *Lactococcus lactis*; Lantibiotic inducible promoters, such as P_{nisA} or P_{nisF} from *Lactococcus lactis* or P_{spaB} from *Bacillus subtilis*; and Galactose-inducible and Thiostrepton-inducible promoters from *Streptomyces lividans*

5 In the absence of inducer for an inducible promoter (e.g., xylose for P_{xylA} ; IPTG for P_{spac} , etc.), low to negligible levels of RAT will be expressed. Thus, when glutamyl tRNA synthetase (ERS) mischarges tRNA^{Gln} with Glu, the cell will no longer be able to convert the Glu-tRNA^{Gln} to Gln-tRNA^{Gln} by transamidation. In mammalian chloroplasts misacylated Glu-tRNA^{Gln} has been shown to be rejected by EF-Tu, and is thus not brought
10 to the ribosome and hence not utilized in protein synthesis (Stanzel-M et al. 1994 Eur. J. Biochem. 219, 435-439). As the cell now has no means of incorporating the amino acid glutamine into nascent proteins, translation will be aborted, leading to cell death. It is provided by the methods of the invention that an analogous will occur in bacteria.

In order to overcome the toxicity caused by ERS, a certain minimum level of RAT
15 enzyme must be present in the cell. This level will relate to a certain level of inducer (herein "level 1"). By adding excess inducer, the levels of RAT will exceed the minimum levels required to relieve ERS toxicity (herein "level 2"). While not wanted to be limited by a mechanism, a schematic of this scenario is illustrated diagrammatically in Figure 2.

In a preferred embodiment, an antimicrobial compound screen may be run at both
20 level 1 and level 2, and hits will be determined, for example, by their ability to cause cell death, as measured by a reduction in OD at 600nm, mimicing the effect of having no inducer of RAT expression present. These hits will include both RAT-specific inhibitors and general bactericidals. The following strategy may be used to differentiate between the two.

25 Hits that cause toxicity at RAT level 1 but not level 2 will be deemed to be RAT-specific inhibitors and not general bactericidals, on the basis that they are not potent enough to inhibit all of the excess of RAT present at level 2, and that general bactericidals will inhibit at both levels. However, not all hits that inhibit at both levels will be general bactericidals. RAT inhibitors that are particularly potent will work at both levels.
30 Therefore, a further screen may be employed for hits in that category. This will involve rerunning the screen using a reduced concentration of these hits, to look for any that only cause toxicity at RAT level 1. These will be deemed to be RAT-specific inhibitors.

An alternative preferred format of this screen will use a different, more sensitive readout to OD reduction in order to assess hits. This will involve expressing luciferase (from the *luxAB* genes) in the cells in the presence of its octanal substrate, which will result in light production. This can be measured using a luminometer. Hits will be identified by their ability to reduce the light output.

Moreover, as *E.coli* belongs to the gamma subdivision of purple bacteria, it does not possess a RAT enzyme. Although neither of the two tRNA(Gln) species in *E.coli* is recognized by *Escherichia coli* (herein *E. coli*) ERS, one of them is recognised by glutamyl tRNA synthetase from *Bacillus subtilis* (herein *B. subtilis*). Applicants used these observations and others to form the basis for the present invention, since Applicants believe that the glutamyl tRNA synthetase enzyme from *S.aureus* will also misacylate the same tRNA(Gln) from *E.coli*; this seems likely as *S.aureus* has RAT activity and *S.aureus* glutamyl tRNA synthetase gene was difficult to clone in *E.coli*.

A method of screening for antimicrobial drugs comprising the steps of: providing at least one cell naturally lacking a RAT gene and comprising at least one recombinant glutamyl tRNA synthetase gene and at least one recombinant RAT gene; contacting the cell with at least one candidate compound; and detecting altered metabolism in the cell of the contacting step. As used herein "RAT gene" means any or all of the genes encoding RAT protein subunits in an enzymatically active or biologically functional RAT protein complex. As used herein "RAT protein" means any or all of the RAT protein subunits in an enzymatically active or biologically functional RAT protein complex.

One preferred embodiment of the invention provides the steps of: expressing *S.aureus* glutamyl tRNA synthetase and RAT, or a variant thereof, in *E.coli*. It is preferred that these gene be on separate plasmids configured such that RAT levels may be regulated. *E. coli* may be contacted with candidate compounds to determine whether they possess antimicrobial activity, and such antimicrobial activity of the candidate compound may be tested. Without being limited by a mechanism of action for the assays of the invention, it is believed that this *E. coli* should not be harmed by the misacylated Glu-tRNA(Gln) as it should be corrected by RAT. Inhibition of RAT activity by a candidate compound possessing antimicrobial activity will result in accumulation of Glu-tRNA(Gln) and consequent cell death. This method has an advantage of being specific for *S.aureus* RAT in a cellular environment and selecting for compounds that display a capacity to penetrate to

the cytosol; achieving compound penetration into *E.coli* is likely to be more demanding than penetration into, e.g. *S.aureus*.

The determining step of the invention may be carried out in many ways in view of the teachings of the present invention. In the methods of the invention, the determining
5 step may be performed using any method to detect altered translation or mysacylation. For example, altered translation is monitored by the introduction of a label into the amino acids of the protein. As a further example, the skilled artisan may detect altered translation directly, such as at the protein level, or indirectly, such as by detecting alterations in the activity of the protein.

10 Another embodiment of the determining step provides detecting Gln-tRNA(Gln) starvation in the host cell following contacting such cell with at least one candidate compound.

Methods are provided herein wherein the determining step comprises detecting or measuring any aspect of altered cellular metabolism, such as detecting or measuring the
15 inhibition of RAT protein activity or RAT gene expression. The determining step may further comprises detecting a toxic accumulation of Glu-tRNA(Gln) and/or cell death.

Following contacting the host cell with at least one candidate compound, if it is determined that there is altered translation, Gln-tRNA(Gln) starvation, cell death, or any other alteration of host cell metabolism detected as provided herein, the candidate
20 compound may be useful as an antimicrobial compound. This may be readily determined using any of the many well known methods for testing antimicrobial activity, such as, for example, by disk diffusion assay followed by an MIC determination.

Preferred method comprise recombinant glutamyl tRNA synthetase and RAT genes that are on an episomal element or integrated into a chromosome of the host cell. It is
25 particularly preferred that in such methods that RAT gene expression level is regulated or regulatable.

The RAT gene, or variants thereof, in the compositions and methods of the invention may be obtained from any source. Methods of the invention may comprise a RAT gene selected from the group consisting of a Gram positive bacterium, a Gram
30 negative bacterium, a streptococcus, *S. pneumoniae*, a staphylococcus, *S. aureus*, enterococci, *Enterococcus faecalis*, *Enterococcus faecium*, a *Bacillus*, and *Bacillus subtilis*, among other bacteria.

Compositions and methods of the invention may comprise glutamyl tRNA synthetase gene, or variants thereof, selected from the group consisting of a Gram positive

bacterium, a Gram negative bacterium, a streptococcus, *S. pneumoniae*, a staphylococcus, *S. aureus*, enterococci, *Enterococcus faecalis*, *Enterococcus faecium*, a *Bacillus*, and *Bacillus subtilis*, among other bacteria.

5 Compositions and methods of the invention may a cell possessing a glutaminyl tRNA synthetase or lacking a RAT gene. Also provided by the invention in an isolated bacterial cell lacking a RAT gene and/or comprising at least one recombinant bacterial glutamyl tRNA synthetase gene and at least one recombinant bacterial RAT gene. These glutamyl tRNA synthetase and/or RAT genes may be present on episomal element or integrated into a chromosome of the host cell. Host cells also provided herein comprise a
10 glutamyl tRNA synthetase gene from *S. aureus*. Host cells are provided possessing a glutaminyl tRNA synthetase and/or lacking a RAT gene.

A preferred embodiment provides a method wherein the polypeptide translated in the screening assay is a nascent polypeptide chain or a substantially purified polypeptide. In any case, the polypeptide in the assay may be labeled using any of the well known
15 polypeptide labeling methods, such as, radiolabeling or chromogenic labeling, or detected using an appropriate antibody and immunoprecipitation reaction. It is preferred that the label used be readily observable or detectable, such as by being luminescent, radiolabeled, colored or fluorescent.

Another preferred embodiment provides a method wherein a test protein is isolated,
20 such as being prepared to a substantially purified form. Test proteins may be isolated using methods well known in the art, including, for example, density centrifugation or gel electrophoresis.

Yet another preferred embodiment provides a method wherein a RAT gene or protein is selected from the group consisting of a Gram positive bacterium, a Gram
25 negative bacterium, a streptococcus, *S. pneumoniae*, a staphylococcus, *S. aureus*, enterococci, *Enterococcus faecalis*, *Enterococcus faecium*, a *Bacillus*, and *Bacillus subtilis*.

A still more preferred embodiment provides a method wherein a RAT gene or protein is derived or isolated from *S. aureus*. Examples of RAT genes and proteins derived or isolated from *S. aureus* are provided in Table 1.

TABLE 1
RAT genes and proteins

Polynucleotide sequence of ratA [SEQ ID NO:1]

5' -
 5 ATGAGCATTTCGCTACGAATCGGTTGAGAATTTATTAACTTTAATAAAAGACAAAAAATCAAACCATCTG
 ATGTTGTTAAAGATATATATGATGCAATTGAAGAGACTGATCCAACAATTAAGTCTTTTCTAGCGCTGGA
 TAAAGAAAAATGCAATCAAAAAAGCGCAAGAATTGGATGAATTACAAGCAAAAGATCAAATGGATGGCAAA
 TTATTTGGTATTCCAATGGGTATAAAAGATAACATTATTACAAACGGATTAGAAACAACATGTGCAAGTA
 10 AAATGTTAGAAGGTTTTGTGCCAATTTACGAATCTACTGTAATGGAAAACTACATAAAGAGAATGCCGT
 TTTAATCGGTAAATTAATATGGATGAGTTTGCAATGGGTGGTTCAACAGAAACATCTTATTTCAAAAA
 ACAGTTAACCCATTTGACCATAAAGCAGTACCAGGTGGTTCATCAGGTGGATCTGCAGCAGCAGTTGCAG
 CTGGCTTAGTACCATTTAGCTTAGGTTTCAGACACAGGTGGTTCATTAGACAACCGGCTGCATATTGTGG
 CGTTGTCGGTATGAAACCAACATACGGTCGTGTATCTCGATTTGGATTAGTTGCTTTTGCATCTTCATTA
 15 GACCAAAATTGGTCCATTGACTCGAAATGTAAAAGATAATGCAATCGTATTAGAAGCTATTTCTGGTGCAG
 ATGTTAATGACTCTACAAGTGCACCAAGTTGATGATGTAGACTTTACATCTGAAATTGGTAAAGATATTAA
 AGGATTAAAAGTTGCATTACCTAAAGAATACTTAGGTGAAGGTGTAGCTGATGACGTAAAAGAAGCAGTT
 CAAAACGCTGTAGAACTTTAAATCTTTAGGTGCTGTCGTTGAGGAAGTATCATTGCCAAATACTAAAT
 TTGGTATTCCATCATATTACGTGATTGCATCATCAGAAGCTTCGTCAAACCTTTCTCGTTTTCACGGAAT
 20 TCGTTATGGTTATCATTCTAAAGAAGCTCATTCAATTAGAAGAATTATATAAAATGTCAAGATCTGAAGGT
 TTCGGTAAAGAAGTAAACGTCGTATTTTCTTAGGTACATTTCGATTAAGTTCAGGTTACTACGATGCTT
 ACTATAAAAAATCTCAAAAAGTTAGAACATTGATTAAAAATGACTTTGATAAAGTATTCGAAAATTATGA
 TGTAGTAGTTGGTCCAACAGCGCCTACAACGCGTTTAATTTAGGTGAAGAAATTGATGATCCATTAACA
 ATGTATGCCAATGATTTATTAAACACCAGTAAACTTAGCTGGATTACCTGGTATTTCTGTTCTTGTG
 25 GACAATCAATGGCCGACCAATCGGTTTACAGTTCATTGGTAAACCATTTCGATGAAAAACGTTATATCG
 TGTGCTTATCAATATGAAACACAATACAATTTACATGACGTTTATGAAAAATTA-3'

Polypeptide sequence of ratA [SEQ ID NO:2] deduced from the sequence of SEQ ID NO:1

NH₂-
 30 MSIRYESVENLLTLIKDKIKPSDVVKDIYDAIEETDPTIKSFLALDKENAIKKAQELDELQAKDQMDGK
 LFGIPMGIKDNIITNGLETTTCASKMLEGFVPIYESTVMEKLHKENAVLIGKLNMDDEFAMGGSTETSIFYKK
 TVNPFDHKAVPGGSSGSAVAAGLVPPSLGSDTGGSSIRQPAAYCGVVGMKPTYGRVSRFGLVAFASSL
 DQIGPLTRNVKDNAIVLEAISGADVNDSTSAPVDDVFTSEIGKDIKGLKVALPKEYLGEVADDVKEAV
 QNAVETLKSLGAVVEEVSLPNTKFGIPSYVVIASSEASSNLSRFDGIRYGYHSKEAHSLEELYKMSRSEG
 35 FGKEVKRRIFLGTFFALSSGYDAYYKKSQKVRTLIKNDFDKVFENYDVVVGPTAPTAFNLGEEIDDPLT
 MYANDLLTTPVNLAGLPGISVPCGQSNRPIGLQFIGKPFDEKTLYRVAYQYETQYNLHDVYEKL-COOH

Polynucleotide sequence of rat B [SEQ ID NO:3]

5' -
 40 ATGCATTTTGAAACAGTTATAGGACTTGAAGTTCACGTAGAGTTAAAAACGGACTCAAAAATGTTTTCTC
 CATCACCAGCGCATTTTGGAGCAGAACCTAACTCAAAATACAAATGTTATCGACTTAGCATATCCAGGTGT
 CTTACCAGTTGTTAATAAGCGTGCAGTAGACTGGGCAATGCGTGCTGCAATGGCACTAAATATGGAAATC

GCAACAGAATCTAAGTTTGACCGTAAGAAC TATTCTATCCAGATAATCCAAAAGCATATCAAATTTCTC
 AATTTGATCAACCAATTTGGTGAAAATGGATATATCGATATCGAAGTCGACGGTGAAACAAAACGAATCGG
 TATTACTCGTCTTCACATGGAAGAAGATGCTGGTAAAGTCAACACATAAAGGTGAGTATTCATTAGTTGAC
 TTGAACCGTCAAGGTACACCGCTAATTGAAATCGTATCTGAACCAGATATTCGTTACCTAAAGAAGCAT
 5 ATGCATATTTAGAAAAATTACGTTCAATTATTC AATACACTGGTGTATCAGACGTTAAGATGGAAGAGGG
 ATCTTTACGTTGTGATGCTAACATCTCTTTGCGTCCATATGGTCAAGAAAAATTTGGTACTAAAGCCGAA
 TTGAAAACTTAAACTCATTTAACTATGTACGTAAAGGTTTAGAATATGAAGAAAAACGCCAAGAAGAAG
 AATTGTTAAATGGTGGAGAAATCGGACAAGAAACACGTCGATTTGATGAATCTACAGGTAAACAATTTT
 AATGCGTGTTAAAGAAGGTTCTGATGATTACCGTTACTTCCCAGAGCCTGACATTGTACCTTTATATATT
 10 GATGATGCTTGGAAAGAGCGTGTTCGTCAGACAATTCCTGAATTACCAGATGAGCGTAAGGCTAAGTATG
 TAAATGAATTAGGTTTACCTGCATACGATGCACACGTATTAACATTGACTAAAGAAATGTCAGATTTCTT
 TGAATCAACAATTGAACACGGTGCAGATGTTAAATTAACATCTAACTGGTTAATGGGTGGCGTAAACGAA
 TATTTAAATAAAAATCAAGTAGAATTATTAGATACTAAATTAACACCAGAAAAATTTAGCAGGTATGATTA
 AACTTATCGAAGACGGAACAATGAGCAGTAAATTCGGAAGAAAGTCTTCCCAGAGTTAGCAGCTAAAGG
 15 TGGTAATGCTAAACAGATTATGGAAGATAATGGCTTAGTTCAAATTTCTGATGAAGCAACACTTCTAAAA
 TTTGTAAATGAAGCATTAGACAATAACGAACAATCAGTTGAAGATTACAAAAATGGTAAAGGCAAAGCTA
 TGGGCTTCTTAGTTGGTCAAATTATGAAAGCGTCTAAAGGTCAAGCTAATCCACAATTAGTAAATCAACT
 ATTTAAACAAGAATTAGATAAAAGA-3'

20 Polypeptide sequence of rat B [SEQ ID NO:4] deduced from the sequence of SEQ ID NO:3.

NH₂-
 MHFETVIGLEVHVELKTD SKMFSPPAHFGAEPNSNTNVIDLAYPGVLPVVKRAVDWAMRAAMALNMEI
 ATESKFDRKNYFYPDNPKAYQISQFDQPIGENGYIDIEVDGETKRIGITRLHMEEDAGKSTHKGEYSLVD
 LNRQGTPLIEIVSEPDIRSPKEAYAYLEKLRSIIQYTGVS DVKMEEGSLRCDANISLRPYGQEKFGTKAE
 25 LKNLNSFN YVRKGLEYEEKRQEEELLNGGEIGQETRRFDESTGKTIILMRVKEGSDDYRYFPEPDIVPLYI
 DDAWKERVQTIPELPDERKAKYVNELGLPAYDAHVLTLTKEMSDFESTIEHGADVKLTSNWL MGGVNE
 YLNKNQVELLDTKLTPENLAGMIKLIEDGTMSSKI AKKVPPELAAKGNAKQIMEDNGLVQISDEATLLK
 FVNEALDNNEQSVEDYKNGKGKAMGFLVGQIMKASKGQANPQLVNQLLKQELDKR-COOH

30 Sequences from *Staphylococcus aureus* ratC polynucleotide sequence [SEQ ID NO:5].

5'-
 ATGACAAAAGTAACACGTGAAGAAGTTGAGCATATCGCGAATCTTGCAAGACTTCAAATTTCTCCTGAAG
 AAACGGAAGAAATGGCCAACACATTAGAAAGCATTTTAGATTTTGCAAAACAAAATGATAGCGCTGATAC
 AGAAGGCGTTGAACCTACATATCACGTTTTAGATTTACAAAACGTTTTACGTGAAGATAAAGCAATTA
 35 GGTATTCGCAAGAATTAGCTTTGAAAAATGCCAAAGAAACAGAAGATGGACAATTTAAAGTGCCCTACAA
 TCATGAATGAGGAGGACGCG-3'

ratC polypeptide sequence [SEQ ID NO:6] deduced from the polynucleotide sequence in this table [SEQ ID NO:5].

NH₂ -

5 MTKVTREEVEHIANLARLQISPEETEEMANTLESILDFAKQNSADTEGVEPTYHVLDLQNVLRDKAIKG
IPQELALKNAKETEDGQFKVPTIMNEEDA-COOH

Any organism possessing a glutamyl-tRNA synthetase activity or not possessing a RAT enzyme, such the gamma subdivision of purple bacteria may be engineered to be useful with certain methods of the invention using a heterogeneous RAT gene and glutamyl-tRNA synthetase (ERS) transformed into such organism. For example, organisms or host cells useful in the methods of the invention include organisms and host cells selected from the group consisting of a eubacterium, an archaeobacterium, gamma subdivision of the purple bacteria, a eukaryote, including lower eukaryotes, such as fungi, protozoa, and cells from higher eukaryotes, such as mammalian cells, CHO, COS, HeLa, C127, 3T3, BHK, 293 and Bowes melanoma cells.

Preferred organisms useful as host cells in the methods of the invention include those which do not possess a RAT enzyme and for which a tRNA(Gln) from such organism is recognised by a heterologous glutamyl tRNA synthetase, such as a glutamyl tRNA synthetase from another species. Applicants used these observations

Once the composition or cells comprising a RAT gene as described in the forgoing is prepared, one or more candidate compounds are added to the composition to test whether any of the candidate compounds is associated with altered translation. If a mixture of candidate compounds is associated with an alteration of RAT expression or activity or glutamyl tRNA synthetase expression or activity, the mixture may be deconvoluted to determine which compound or compounds is active. One method to achieve this is to test each component of the mixture separately in the assay. Deconvolution may also be performed using any of the known deconvolution methods.

It is preferred that the method of the invention is formatted for high throughput screening (herein "HTS"). Skilled artisans can readily adapt the method of the invention for HTS. A particularly preferred embodiment of the screening methods of the invention is a high throughput screen for compounds that interfere with the proper functioning of RAT gene expression or protein and/or glutamyl tRNA synthetase gene expression or protein, such as compounds that are associated with altered translation.

Potential antimicrobial compounds identified using the method of the invention include, among other things, small organic molecules, polynucleotides, peptides, polypeptides and antibodies that bind RAT polynucleotides or polypeptides, or mimic the activity of a RAT polypeptides.

5 Potential antagonists include a small molecule that binds to RAT polynucleotides or polypeptides thereby preventing binding of natural factors, such as, for example, tRNAs, such that normal biological activity is prevented. Examples of small molecules include but are not limited to small organic molecules, peptides or peptide-like molecules.

10 The invention further provides assay packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Examples of preferred kits are kits comprising at least one bacterial cell lacking a RAT gene and the cell comprising at least one bacterial glutamyl tRNA synthetase gene and at least one bacterial RAT gene. A further preferred kit comprises a polynucleotide encoding a RAT gene and a polynucleotide encoding a glutamyl tRNA
15 synthetase gene. Kits comprising a RAT gene expressibly linked to an inducible promoter are also preferred.

Each reference disclosed herein is incorporated by reference herein in its entirety. Any patent application to which this application claims priority is also incorporated by
20 reference herein in its entirety.

What is claimed is:

1. A method of screening for antimicrobial drugs comprising the steps of:
providing at least one cell naturally lacking a RAT gene and comprising at least one
5 recombinant or wild type glutamyl tRNA synthetase gene and at least one recombinant
RAT gene; contacting said cell with at least one candidate compound; and detecting altered
metabolism in said cell of the contacting step.
2. The method of claim 1 wherein the recombinant glutamyl tRNA
synthetase and RAT genes are on an episomal element or integrated into a chromosome of
10 said cell.
3. The method of claim 2 wherein RAT gene expression level is regulated.
4. The method of claim 1 wherein at least one RAT gene is a RAT gene
selected from the group consisting of a Gram positive bacterium, a Gram negative
bacterium, a streptococcus, *S. pneumoniae*, a staphylococcus, *S. aureus*, enterococci,
15 *Enterococcus faecalis*, *Enterococcus faecium*, a Bacillus, and *Bacillus subtilis*.
5. The method of claim 1 wherein the glutamyl tRNA synthetase gene is a
glutamyl tRNA synthetase gene selected from the group consisting of a Gram positive
bacterium, a Gram negative bacterium, a streptococcus, *S. pneumoniae*, a staphylococcus, *S.*
aureus, enterococci, *Enterococcus faecalis*, *Enterococcus faecium*, a Bacillus, and *Bacillus*
20 *subtilis*.
6. The method of claim 1 wherein said altered metabolism comprises
inhibition of RAT protein activity.
7. The method of claim 1 wherein said detecting step further comprises
detecting a toxic accumulation of Glu-tRNA(Gln) or toxic incorporation of a glutamyl
25 residue in place of a glutaminyl residue in nascent protein chains.
8. The method of claim 1 wherein said detecting step further comprises
detecting cell death or a reduction in growth rate or amount.
9. The method of claim 1 wherein said cell possesses or lacks a glutaminyl
tRNA synthetase or possesses or lacks a RAT gene.
- 30 10. An isolated bacterial cell lacking a RAT gene and comprising at least one
recombinant or wild type bacterial glutamyl tRNA synthetase gene and at least one
recombinant bacterial RAT gene.
11. The cell of claim 10 wherein said glutamyl tRNA synthetase and RAT
genes are on episomal element or integrated into a chromosome of said cell.

12. The cell of claim 10 wherein said RAT gene expression level is regulated.
13. The cell of claim 10 wherein said RAT gene is a *S. aureus* RAT gene.
14. The cell of claim 10 wherein said glutamyl tRNA synthetase gene is a *S. aureus* glutamyl tRNA synthetase gene.
- 5 15. The cell of claim 10 wherein said cell possesses a glutamyl tRNA synthetase or lacks a RAT gene.
16. The method of claim 1 wherein said detecting step further comprises detecting altered translation.
17. The method of claim 1 wherein said detecting step further comprises
10 detecting altered test protein.
18. A kit comprising at least one bacterial cell lacking a RAT gene and said cell comprising at least one bacterial glutamyl tRNA synthetase gene and at least one bacterial RAT gene.
19. A kit comprising a polynucleotide encoding a RAT gene and a
15 polynucleotide encoding a glutamyl tRNA synthetase gene.
20. A polynucleotide comprising a RAT expressibly linked to an inducible promoter.




FIG. 1




FIG. 2




FIG. 3

SEQUENCE LISTING

<110> Michael T. Black
Howard Kallender

<120> RAT WHOLE CELL ASSAY

<130> GM50039

<140> Unknown

<141> October 2, 1998

<160> 6

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1455

<212> DNA

<213> *Staphylococcus aureus*

<400> 1

atgagcattc gctacgaatc ggttgagaat ttattaactt taataaaaga caaaaaaatc	60
aaaccatctg atgttggttaa agatatatat gatgcaattg aagagactga tccaacaatt	120
aagtcttttc tagcgctgga taaagaaaat gcaatcaaaa aagcgcaaga attggatgaa	180
ttacaagcaa aagatcaaat ggatggcaaa ttatttggtt ttccaatggg tataaaagat	240
aacattatta caaacggatt agaaacaaca tgtgcaagta aaatggttaga aggttttgtg	300
ccaattttacg aatctactgt aatggaaaaa ctacataaag agaatgccgt tttaatcggg	360
aaattaaata tggatgagtt tgcaatgggt ggttcaacag aaacatctta tttcaaaaaa	420
acagttaacc catttgacca taaagcagta ccaggtggtt catcaggtgg atctgcagca	480
gcagttgcag ctggcttagt accatttagc ttaggttcag acacaggtgg ttcaattaga	540
caaccggctg catattgtgg cgttgctcgg atgaaaccaa catacggctg tgtatctcga	600
tttgatttag ttgcttttgc atcttcatta gaccaaattg gtccattgac tcgaaatgta	660
aaagataatg caatcgtatt agaagctatt tctggtgcag atgttaatga ctctacaagt	720
gcaccagttg atgatgtaga ctttacatct gaaattggta aagatattaa aggattaaaa	780
gttgcatcac cttaaagaata cttaggtgaa ggtgtagctg atgacgtaaa agaagcagtt	840
caaaacgctg tagaaacttt aaaatcttta ggtgctgtcg ttgaggaagt atcattgcc	900
aataactaaat ttggtattcc atcatattac gtgattgcat catcagaagc ttcgtcaaac	960
ctttctcgtt ttgacggaat tcgttatggt tatcattcta aagaagctca ttcattagaa	1020
gaattatata aaatgtcaag atctgaaggt ttcggtaaag aagtaaaacg tcgtattttc	1080
ttaggtagat ttgcattaag ttcagggttac tacgatgctt actataaaaa atctcaaaaa	1140
gttagaacat tgattaaaaa tgactttgat aaagtattcg aaaattatga tgtagtagtt	1200
ggtccaacag gcctacaac tgcgtttaat ttaggtgaag aaattgatga tccattaaca	1260

- 2 -

atgtatgccca atgatttatt aacaacacca gtaaacttag ctggattacc tggattttct 1320
 gttccttgtg gacaatcaaa tggccgacca atcggtttac agttcattgg taaaccattc 1380
 gatgaaaaaa cgttatatcg tgtcgcttat caatatgaaa cacaatacaa tttacatgac 1440
 gtttatgaaa aatta 1455

<210> 2

<211> 485

<212> PRT

<213> *Staphylococcus aureus*

<400> 2

Met Ser Ile Arg Tyr Glu Ser Val Glu Asn Leu Leu Thr Leu Ile Lys
 1 5 10 15
 Asp Lys Lys Ile Lys Pro Ser Asp Val Val Lys Asp Ile Tyr Asp Ala
 20 25 30
 Ile Glu Glu Thr Asp Pro Thr Ile Lys Ser Phe Leu Ala Leu Asp Lys
 35 40 45
 Glu Asn Ala Ile Lys Lys Ala Gln Glu Leu Asp Glu Leu Gln Ala Lys
 50 55 60
 Asp Gln Met Asp Gly Lys Leu Phe Gly Ile Pro Met Gly Ile Lys Asp
 65 70 75 80
 Asn Ile Ile Thr Asn Gly Leu Glu Thr Thr Cys Ala Ser Lys Met Leu
 85 90 95
 Glu Gly Phe Val Pro Ile Tyr Glu Ser Thr Val Met Glu Lys Leu His
 100 105 110
 Lys Glu Asn Ala Val Leu Ile Gly Lys Leu Asn Met Asp Glu Phe Ala
 115 120 125
 Met Gly Gly Ser Thr Glu Thr Ser Tyr Phe Lys Lys Thr Val Asn Pro
 130 135 140
 Phe Asp His Lys Ala Val Pro Gly Gly Ser Ser Gly Gly Ser Ala Ala
 145 150 155 160
 Ala Val Ala Ala Gly Leu Val Pro Phe Ser Leu Gly Ser Asp Thr Gly
 165 170 175
 Gly Ser Ile Arg Gln Pro Ala Ala Tyr Cys Gly Val Val Gly Met Lys
 180 185 190
 Pro Thr Tyr Gly Arg Val Ser Arg Phe Gly Leu Val Ala Phe Ala Ser
 195 200 205
 Ser Leu Asp Gln Ile Gly Pro Leu Thr Arg Asn Val Lys Asp Asn Ala
 210 215 220
 Ile Val Leu Glu Ala Ile Ser Gly Ala Asp Val Asn Asp Ser Thr Ser
 225 230 235 240
 Ala Pro Val Asp Asp Val Asp Phe Thr Ser Glu Ile Gly Lys Asp Ile
 245 250 255

- 3 -

Lys Gly Leu Lys Val Ala Leu Pro Lys Glu Tyr Leu Gly Glu Gly Val
 260 265 270
 Ala Asp Asp Val Lys Glu Ala Val Gln Asn Ala Val Glu Thr Leu Lys
 275 280 285
 Ser Leu Gly Ala Val Val Glu Glu Val Ser Leu Pro Asn Thr Lys Phe
 290 295 300
 Gly Ile Pro Ser Tyr Tyr Val Ile Ala Ser Ser Glu Ala Ser Ser Asn
 305 310 315 320
 Leu Ser Arg Phe Asp Gly Ile Arg Tyr Gly Tyr His Ser Lys Glu Ala
 325 330 335
 His Ser Leu Glu Glu Leu Tyr Lys Met Ser Arg Ser Glu Gly Phe Gly
 340 345 350
 Lys Glu Val Lys Arg Arg Ile Phe Leu Gly Thr Phe Ala Leu Ser Ser
 355 360 365
 Gly Tyr Tyr Asp Ala Tyr Tyr Lys Lys Ser Gln Lys Val Arg Thr Leu
 370 375 380
 Ile Lys Asn Asp Phe Asp Lys Val Phe Glu Asn Tyr Asp Val Val Val
 385 390 395 400
 Gly Pro Thr Ala Pro Thr Thr Ala Phe Asn Leu Gly Glu Glu Ile Asp
 405 410 415
 Asp Pro Leu Thr Met Tyr Ala Asn Asp Leu Leu Thr Thr Pro Val Asn
 420 425 430
 Leu Ala Gly Leu Pro Gly Ile Ser Val Pro Cys Gly Gln Ser Asn Gly
 435 440 445
 Arg Pro Ile Gly Leu Gln Phe Ile Gly Lys Pro Phe Asp Glu Lys Thr
 450 455 460
 Leu Tyr Arg Val Ala Tyr Gln Tyr Glu Thr Gln Tyr Asn Leu His Asp
 465 470 475 480
 Val Tyr Glu Lys Leu
 485

<210> 3

<211> 1425

<212> DNA

<213> *Staphylococcus aureus*

<400> 3

atgcattttg aaacagttat aggacttgaa gttcacgtag agttaaaaac ggactcaaaa 60
 atgttttctc catcaccagc gcattttgga gcagaacctt actcaaatac aaatgtttatc 120
 gacttagcat atccaggtgt cttaccagtt gttaataagc gtgcagtaga ctgggcaatg 180
 cgtgctgcaa tggcactaaa tatggaaatc gcaacagaat ctaagtttga ccgtaagaac 240
 tattttctatc cagataatcc aaaagcatat caaatttctc aatttgatca accaattggt 300
 gaaaatggat atatcgatat cgaagtgcac ggtgaaacaa aacgaatcgg tattactcgt 360
 cttcacatgg aagaagatgc tggttaagtca acacataaag gtgagtattc attagttgac 420

- 4 -

```

ttgaaccgtc aaggtaacc gctaattgaa atcgtatctg aaccagatat tcgttcacct 480
aaagaagcat atgcatattt agaaaaatta cgttcaatta ttcaatacac tgggtgatca 540
gacgttaaga tggaagaggg atctttacgt tgtgatgcta acatctcttt gcgtccatat 600
ggtcaagaaa aatttggtac taaagccgaa ttgaaaaact taaactcatt taactatgta 660
cgtaaagggt tagaatatga agaaaaacgc caagaagaag aattgttaaa tgggtggagaa 720
atcggacaag aaacacgtcg atttgatgaa tctacaggta aaacaatttt aatgcgtggt 780
aaagaagggt ctgatgatta ccgttacttc ccagagcctg acattgtacc tttatatatt 840
gatgatgctt ggaaagagcg tgttcgtcag acaattcctg aattaccaga tgagcgtaag 900
gctaagtatg taaatgaatt aggtttacct gcatacgtg cacacgtatt aacattgact 960
aaagaaatgt cagatttctt tgaatcaaca attgaacacg gtgcagatgt taaattaaca 1020
tctaactggt taatgggtgg cgtaaacgaa tatttaaata aaaatcaagt agaattatta 1080
gatactaaat taacaccaga aaatttagca ggtatgatta aacttatcga agacggaaca 1140
atgagcagta aaattgcgaa gaaagtcttc ccagagttag cagctaaagg tggtaatgct 1200
aaacagatta tggaagataa tggcttagtt caaatttctg atgaagcaac acttctaaaa 1260
tttgtaaata aagcattaga caataacgaa caatcagttg aagattacaa aaatggtaaa 1320
ggcaaagcta tgggcttctt agttggtcaa attatgaaag cgtctaaagg tcaagctaata 1380
ccacaattag taaatcaact attaaaacaa gaattagata aaaga 1425

```

<210> 4

<211> 475

<212> PRT

<213> *Staphylococcus aureus*

<400> 4

```

Met His Phe Glu Thr Val Ile Gly Leu Glu Val His Val Glu Leu Lys
1           5           10           15
Thr Asp Ser Lys Met Phe Ser Pro Ser Pro Ala His Phe Gly Ala Glu
          20           25           30
Pro Asn Ser Asn Thr Asn Val Ile Asp Leu Ala Tyr Pro Gly Val Leu
          35           40           45
Pro Val Val Asn Lys Arg Ala Val Asp Trp Ala Met Arg Ala Ala Met
          50           55           60
Ala Leu Asn Met Glu Ile Ala Thr Glu Ser Lys Phe Asp Arg Lys Asn
65           70           75           80
Tyr Phe Tyr Pro Asp Asn Pro Lys Ala Tyr Gln Ile Ser Gln Phe Asp
          85           90           95
Gln Pro Ile Gly Glu Asn Gly Tyr Ile Asp Ile Glu Val Asp Gly Glu
          100          105          110
Thr Lys Arg Ile Gly Ile Thr Arg Leu His Met Glu Glu Asp Ala Gly
          115          120          125
Lys Ser Thr His Lys Gly Glu Tyr Ser Leu Val Asp Leu Asn Arg Gln
          130          135          140
Gly Thr Pro Leu Ile Glu Ile Val Ser Glu Pro Asp Ile Arg Ser Pro
145          150          155          160

```

- 5 -

Lys Glu Ala Tyr Ala Tyr Leu Glu Lys Leu Arg Ser Ile Ile Gln Tyr
 165 170 175
 Thr Gly Val Ser Asp Val Lys Met Glu Glu Gly Ser Leu Arg Cys Asp
 180 185 190
 Ala Asn Ile Ser Leu Arg Pro Tyr Gly Gln Glu Lys Phe Gly Thr Lys
 195 200 205
 Ala Glu Leu Lys Asn Leu Asn Ser Phe Asn Tyr Val Arg Lys Gly Leu
 210 215 220
 Glu Tyr Glu Glu Lys Arg Gln Glu Glu Glu Leu Leu Asn Gly Gly Glu
 225 230 235 240
 Ile Gly Gln Glu Thr Arg Arg Phe Asp Glu Ser Thr Gly Lys Thr Ile
 245 250 255
 Leu Met Arg Val Lys Glu Gly Ser Asp Asp Tyr Arg Tyr Phe Pro Glu
 260 265 270
 Pro Asp Ile Val Pro Leu Tyr Ile Asp Asp Ala Trp Lys Glu Arg Val
 275 280 285
 Arg Gln Thr Ile Pro Glu Leu Pro Asp Glu Arg Lys Ala Lys Tyr Val
 290 295 300
 Asn Glu Leu Gly Leu Pro Ala Tyr Asp Ala His Val Leu Thr Leu Thr
 305 310 315 320
 Lys Glu Met Ser Asp Phe Phe Glu Ser Thr Ile Glu His Gly Ala Asp
 325 330 335
 Val Lys Leu Thr Ser Asn Trp Leu Met Gly Gly Val Asn Glu Tyr Leu
 340 345 350
 Asn Lys Asn Gln Val Glu Leu Leu Asp Thr Lys Leu Thr Pro Glu Asn
 355 360 365
 Leu Ala Gly Met Ile Lys Leu Ile Glu Asp Gly Thr Met Ser Ser Lys
 370 375 380
 Ile Ala Lys Lys Val Phe Pro Glu Leu Ala Ala Lys Gly Gly Asn Ala
 385 390 395 400
 Lys Gln Ile Met Glu Asp Asn Gly Leu Val Gln Ile Ser Asp Glu Ala
 405 410 415
 Thr Leu Leu Lys Phe Val Asn Glu Ala Leu Asp Asn Asn Glu Gln Ser
 420 425 430
 Val Glu Asp Tyr Lys Asn Gly Lys Gly Lys Ala Met Gly Phe Leu Val
 435 440 445
 Gly Gln Ile Met Lys Ala Ser Lys Gly Gln Ala Asn Pro Gln Leu Val
 450 455 460
 Asn Gln Leu Leu Lys Gln Glu Leu Asp Lys Arg
 465 470 475

<210> 5

<211> 300

<212> DNA

- 6 -

<213> *Staphylococcus aureus*

<400> 5

```

atgacaaaag taacacgtga agaagttgag catatcgaga atcttgcaag acttcaaatt      60
tctcctgaag aaacggaaga aatggccaac acattagaaa gcattttaga ttttgcaaaa      120
caaaatgata gcgctgatac agaaggcgtt gaacctacat atcacgtttt agatttacia      180
aacgtttttac gtgaagataa agcaattaaa ggtattccgc aagaattagc tttgaaaaat      240
gccaaagaaa cagaagatgg acaattttaa gtgcctacaa tcatgaatga ggaggacgcg      300

```

<210> 6

<211> 100

<212> PRT

<213> *Staphylococcus aureus*

<400> 6

```

Met Thr Lys Val Thr Arg Glu Glu Val Glu His Ile Ala Asn Leu Ala
 1              5              10              15
Arg Leu Gln Ile Ser Pro Glu Glu Thr Glu Glu Met Ala Asn Thr Leu
      20              25              30
Glu Ser Ile Leu Asp Phe Ala Lys Gln Asn Asp Ser Ala Asp Thr Glu
      35              40              45
Gly Val Glu Pro Thr Tyr His Val Leu Asp Leu Gln Asn Val Leu Arg
      50              55              60
Glu Asp Lys Ala Ile Lys Gly Ile Pro Gln Glu Leu Ala Leu Lys Asn
      65              70              75              80
Ala Lys Glu Thr Glu Asp Gly Gln Phe Lys Val Pro Thr Ile Met Asn
              85              90              95
Glu Glu Asp Ala
              100

```

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/20582**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : C12Q1/68; C12N 15/00

US CL : 435/172.3, 6

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/172.3, 6

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

DIALOG, APS

search terms: rat gene, transferase?, tna, synthetase?, misaminoacylat?, glutamyl?, glu-tna, gln (2a) tna

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,643,722 A (ROTHSCHILD et al) 01 July 1997, see title, abstract and claims, and entire document.	1-20
A	US 5,646,024 A (LEEMANS et al) 08 July 1997, see title, abstract, claims and entire document.	1-20

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

18 DECEMBER 1998

Date of mailing of the international search report

25 JAN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GINNY PORTNER

Telephone No. (703) 308-0196

